# User guide

## Compartmental Analysis (PET)

— LIFEx —

C. Nioche, I. Buvat

LIFEx version 7.4.n, Last update of document: 2023/03/20

Introduction

 $M_{\text{odels}}$ 

# Chapter 1 Introduction

## 1.1 Models

Compartmental analysis in LIFEx application come from of lhsol 2.0.2 program 2002-2012 by Turku PET Centre

Fitting of full or reduced compartmental model to plasma and tissue time-activity curves (TACs) to estimate the model parameters:

Where Model is one of these:

- lhsolK1: K1 (for assuming k2=k3=k4=k5=k6=0)
- lhsolvk1: K1 Vp(%) (for assuming k2=k3=k4=k5=k6=0)
- lhsolk2: K1 k2 K1/k2 (for assuming k3=k4=k5=k6=0)
- lhsolvk2: K1 k2 Vp(%) K1/k2 (for assuming k3=k4=k5=k6=0)

### 1.1 Models

- lhsolk3: K1 k2 k3 K1/k2 Ki (for assuming k4=k5=k6=0)
- lhsolvk3: K1 k2 k3 Vp(%) K1/k2 Ki (for assuming k4=k5=k6=0)
- lhsolk4: K1 k2 k3 k4 K1/k2 k3/k4 Vd (for assuming k5=k6=0)
- lhsolvk4: K1 k2 k3 k4 Vp(%) K1/k2 k3/k4 Vd (for assuming k5=k6=0)

Compartmental models are transformed into general linear least squares functions (1, 2, 3), which are solved using Lawson-Hanson non-negative least squares (NNLS) algorithm (4). Linear parameters are always >=0, but compartmental model parameters may get negative estimates. Note that rate constants and macroparameters are represented per volume (measured by PET) including vascular volume.




Figure 1.1: main screenshot of PT Compartmental protocol

### References:

- 1. Blomqvist G. On the construction of functional maps in positron emission tomography. J Cereb Blood Flow Metab 1984;4:629-632.
- 2. Gjedde A, Wong DF. Modeling neuroreceptor binding of radioligands in vivo. In: Quantitative imaging: neuroreceptors, neurotransmitters, and enzymes. (Eds. Frost JJ, Wagner HM Jr). Raven Press, 1990, 51-79.

- 3. Oikonen V. Multilinear solution for 4-compartment model: I. Tissue compartments in series. http://www.turkupetcentre.net/reports/tpcmod0023.pdf
- 4. Lawson CL & Hanson RJ. Solving least squares problems. Prentice-Hall, 1974.